

GLUT3cKO and PKM1cKO do not alter post-synaptic electrophysiology under standard high-glucose conditions

We next investigated the consequences of PKM1 or GLUT3 deficiency on the function of CA1 pyramidal cells. First, we used hippocampal slices from 20- to 23-month-old mice to interrogate the CA3-CA1 pathway by stimulating Schaffer collaterals and recording the response in both the dendritic layer (str. Radiatum, field postsynaptic potentials [fPSPs]) and the pyramidal cell layer (str. Pyramidale, local field potentials [LFPs]) (Figure S6A). We found that loss of either GLUT3 or PKM1 did not affect the pre-synaptic dynamics, seen as paired-pulse ratio at varying inter-pulse intervals (Figures S6B and S6H), the post-synaptic response to increasing stimulation intensities (Figures S6C and S6I), or cellular excitability (population spike integral vs. fPSP slope relationship; Figures S6D and S6J). Network function in response to repetitive synaptic stimulation (30 s, 10-Hz train) was also unaltered, with knockout (KO) slices exhibiting a robust response comparable to that in WT both in dendritic (Figures S6E and S6K) and cellular (Figures S6F, S6G, S6L, and S6M) layers. These findings suggest that PKM1 or GLUT3 deficiency in CA1 pyramidal cells does not significantly alter their post-synaptic electrophysiological function under standard electrophysiologic conditions, including high glucose concentrations (10 mM), although we did not assess function under physiologic brain glucose levels (≈ 1 –1.5 mM). In addition, these recordings assess the post-synaptic response in CA1 dendritic and somatic layers and therefore would not detect any potential changes downstream at the CA1 pyramidal cells' axonal boutons. The latter requires interrogation of the output of CA1, which is technically challenging in *ex vivo* brain slice preparations.

GLUT3cKO neurons have decreased glucose uptake and levels that are unresponsive to changes in external glucose

Our behavioral studies indicate a requirement for glucose uptake and glycolysis for CA1 neuronal function. Our finding that GLUT3cKO decreased brain size and body weight in female mice, while PKM1cKO did not, also suggests that neurons require glucose uptake for functions besides glycolysis, although differences in the ability to compensate for GLUT3 versus PKM1 loss may also contribute. To gain insight into how GLUT3 influences glucose metabolism in neurons, we examined the impact of GLUT3cKO on glycolytic gene expression in CA1 neurons, but this was unchanged (Figure 6A).

We hypothesized that decreased glucose uptake across physiologically relevant concentrations would decrease the levels of cytosolic glucose and downstream glucose metabolites. We therefore used targeted and untargeted metabolomics to examine the impact of lowering GLUT3 levels with CRISPRi on glucose metabolites. iPSC lines co-expressing an inducible

dCas9-KRAB and either GLUT3 (GLUT3 KD) or an NTG (control) sgRNA were differentiated into neurons, and then cultured for 11 days, at which time GLUT3 KD neurons showed near-complete KD of GLUT3 expression (Figure S7A).

Neurons were then incubated with [U-¹³C]glucose for 24 h prior to harvesting for metabolomics.²³ In control cells, the incorporation of ¹³C-labeled metabolites into downstream glycolytic, TCA, and PPP metabolites decreased with decreasing extracellular [U-¹³C]glucose, spanning concentrations typically found in cell culture and *in vivo* in physiologic and pathologic conditions (0.15–20.5 mM) (Figure S7B).

Interestingly, intracellular glucose was decreased in GLUT3KD neurons incubated with [U-¹³C]glucose for 24 h at the physiological 1.5 mM glucose concentration but not at 20.5 mM glucose (Figure S7B). Specifically, the percentage of glucose labeled by [U-¹³C] decreased from $\approx 60\%$ in the NTG neurons to 30% in the GLUT3 KD neurons, while total levels dropped to $<20\%$ of control values (Figures 6B and S7C), consistent with decreased glucose uptake due to GLUT3 KD. Moreover, GLUT3 KD neurons had markedly decreased labeled fractions of the glycolytic metabolite PEP, as well as the PPP metabolites 6P-gluconate, and ribose 5-phosphate (R5P). There were no changes in either the labeled or total fractions of TCA metabolites. Indeed, despite taking in less [U-¹³C]glucose, neurons with GLUT3 KD still derived an amount of citrate and lactate from [U-¹³C]glucose equivalent to control neurons ($>70\%$ and 50% respectively). This suggests that, when glucose is limiting, neurons prioritize metabolism of glucose metabolites through the TCA cycle and aerobic glycolysis, rather than through the PPP. In contrast, in higher glucose levels typically used for cell culture, GLUT3 KD had little or no impact on either labeled or total levels of glucose or downstream glycolytic, PPP, or TCA metabolites, suggesting that sufficient glucose can enter neurons through other GLUTs under these conditions (Figure S7B).

We next examined relative glucose levels in individual neurons in culture. GLUT3^{lox/lox} hippocampal neurons were co-transfected with a glucose sensor (iGlucoSnFR-mRuby) and either Cre (to delete GLUT3, GLUT3KO) or empty vector (control). We found that GLUT3KO neurons had similar basal glucose levels to controls at the cell body (Figure 6C). However, in contrast to controls, their glucose levels were unresponsive to changes in the extracellular glucose level, consistent with a profound deficit in glucose uptake in the absence of GLUT3 (Figure 6D). We next examined the impact of electrical field stimulation (5 Hz, 5.5 min) to increase the energy requirements of neurons by increasing neural activity.^{30,52,53} At the cell body, when extracellular glucose was the sole energy substrate, glucose levels decreased to a similar extent in control and GLUT3KO neurons (Figure 6E). As expected, the magnitude and speed of drop was somewhat greater when all glucose uptake was blocked

(I) GLUT3KO neurons have similar basal ATP levels at cell bodies (left), and in response to electrical stimulation (5 Hz, 5.5 min) to increase the ATP demand (right). $n = 11$ –13 coverslips/group, two or three cells/coverslip from five independent experiments.

(J) GLUT3KO synaptic boutons have decreased ATP levels (left), and their ATP levels decrease less in response to stimulation (5 Hz, 5.5 min). $n = 6$ –7 coverslips/group, three to five synapses/coverslip from three independent experiments.

ns, not significant; * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$, by unpaired t tests (C, F, I, J) or two-way ANOVA with Tukey's multiple comparison (A and B). Brackets in graphs show significance of linear mixed modeling for the interaction of genotype and time (D, E, G, H, I right, J right) with Sidak's multiple comparison.

(no glucose and cytochalasin B). These results suggest that, at the cell body, GLUT3KO neurons decrease their rate of glucose consumption to match decreased uptake, such that glucose levels decrease at a similar rate to controls.

We next examined the impact of GLUT3KO on glucose levels at the synapse, where GLUT3 may be enriched.⁵⁴ At baseline, GLUT3KO neurons had lower glucose level at the synapse than controls (Figure 6F), and their glucose levels were unresponsive to changes in the extracellular glucose level (Figure 6G). Moreover, in the presence of extracellular glucose, intracellular glucose levels declined far more rapidly and to a greater extent in GLUT3KO neurons than controls in response to electrical field stimulation (Figure 6H, left). This suggests that, at the nerve terminal, GLUT3KO neurons consume glucose rapidly in response to electrical stimulation and are unable to replenish their supply, resulting in a marked decline in glucose levels. Blocking all glucose uptake eventually caused glucose levels in control neurons to drop to GLUT3KO levels (Figure 6H, right).

Neurons require glycolysis to maintain synaptic ATP levels

Our electrophysiology data indicate that CA1 neuronal cell bodies in GLUT3cKO slices have sufficient ATP levels to support increased neural activity. However, these studies were unable to provide insight into ATP levels at the synapse, which may be particularly vulnerable to energy failure^{30,52} because the proportional increase in energy expenditure with neural activity may be greater in synapses than the cell body. To determine if GLUT3KO decreases glucose-derived ATP production in individual neurons, including at the synapse, GLUT3^{lox/lox} and WT neurons were co-transfected with either Cre or empty vector, and an ATP-fluorescence resonance energy transfer (FRET) sensor made of the ϵ subunit of the bacterial FoF₁-ATP synthase (an ATP-binding protein) flanked by mApple and mClover, which is sensitive to physiologic changes in ATP levels.^{55,56} GLUT3KO had no impact on ATP levels at the cell body in basal conditions (30 mM glucose, 10 mM pyruvate), in which cells could derive ATP from both glycolysis and aerobic respiration (Figure 6I). Moreover, GLUT3KO ATP levels dropped similarly to controls with sustained electrical field stimulation (5 Hz, 5.5 min). However, basal ATP levels were lower in GLUT3KO nerve terminals than in controls, and ATP levels in GLUT3KO nerve terminals failed to decline further during sustained electrical stimulation (Figure 6J), suggesting that GLUT3KO neurons either fail to fire in response to the electrical stimulation despite consuming glucose or that they rely on alternative fuel sources to maintain ATP levels. These findings show that neurons rely on glycolysis to maintain ATP levels at the synapse, consistent with the hypothesis that glycolysis is important for rapid response to increased ATP demand to support neural activity.¹³

Compensatory pathways in GLUT3KO neurons

Although GLUT3KO disrupts energy metabolism and neuronal function, neurons can nonetheless survive and metabolize glucose without GLUT3. To delineate potential compensatory mechanisms, we used spatial genomics to examine gene expression in CA1 neurons in 7-month-old GLUT3cKO and WT mice. Targeted examination revealed no compensatory changes in other GLUTs (Figure 7A). There were also no increases in the expression of TCA genes in GLUT3cKO neurons (Figure 7B). To search for other compensatory mechanisms, we performed an unbiased analysis of differentially expressed genes in CA1 neurons. The resulting hits were enriched with genes related to either carbohydrate or mitochondrial energy metabolism (Figures 7C–7E). The mitochondrial genes included *lars2*, which encodes mitochondrial isoleucine-tRNA synthetase and whose mutation produces a range of clinical disorders with impaired respiratory chain function, including Leigh syndrome^{57,58}; *Echdc2* (enoyl-CoA hydratase domain-containing protein 2), a mitochondrial protein involved in beta oxidation of fatty acids and valine catabolism,^{59,60} and whose deficiency can also cause Leigh syndrome⁶¹; and melanocyte inducing transcription factor (*Mitf*), which regulates both PDH⁶² and PGC-1 α ,⁶³ while increasing oxidative phosphorylation,⁶⁴ boosting ATP levels under respiratory conditions²³ and protecting against glucose restriction.⁶⁵ Indeed, *Mitf* expression was increased by immunofluorescence in CA1 neurons from GLUT3cKO mice (Figure S7D).

Although most hits had similar effects on male and female GLUT3KO neurons, we also identified several genes with sex-specific effects (Figures 7D and 7E). In particular, *Pdp2* (pyruvate dehydrogenase phosphatase catalytic subunit 2), which diverts metabolism to oxidative phosphorylation, was lower at baseline in male versus female CA1 neurons, while GLUT3cKO increased *Pdp2* expression to similar levels as those in female WT and GLUT3cKO neurons. Another mitochondrial gene, *Bcs1l* (mitochondrial chaperone BCS1), needed for the assembly of mitochondrial respiratory chain complex III, was decreased in female GLUT3cKO neurons. *Bcs1l* immunofluorescence was also decreased in CA1 neurons from female GLUT3cKO mice (Figure S7D). *Bcs1l* expression may be downregulated in response to an overall decrease in mitochondrial respiration in GLUT3cKO mice, but further investigation is required to delineate the impacts of these gene expression changes on ATP production when glucose is limiting.

Pathway analysis of CA1 gene expression in GLUT3cKO mice also identified an enrichment of galactose metabolism as a potential compensatory mechanism (Table S2). In particular, *Gale* (UDP-galactose-epimerase), which plays a key role in the catabolism of galactose into glucose-6-phosphate such that galactose can enter the glycolytic pathway via the Leloir pathway,⁶⁶ was upregulated in GLUT3cKO mice (Figure 7E). Galactose is

(H–J) Relative amounts and [U-¹³C]galactose fractional labeling of Leloir, PPP, and glycolytic metabolites in GLUT3 KD and NTG neurons after 24 h of culture. n = 4 samples/group.

(K–M) Relative total amounts and [U-¹³C]glucose fractional labeling of Leloir, PPP, and glycolytic pathway metabolites in GLUT3 KD neurons relative to NTG neurons when treated with either 1.5 mM [U-¹³C]glucose or 1.5:1.5 mM [U-¹³C]glucose:galactose for 24 h. n = 4 samples/group.

*p < 0.05, **p < 0.01, ***p < 0.001 by unpaired t test and two-way ANOVA with either Tukey's multiple comparisons test (D, E, H–M) or Sidak's multiple comparisons test (F).

also imported across the cell membrane via several glucose transporters, including GLUT3.⁶⁷ Two genes involved in the metabolism of UDP-glucose demonstrated sex-specific differences with GLUT3cKO: Galt (galactose-1-phosphate uridylyl-transferase) and Ugdh (UDP-glucose 6-dehydrogenase). Expression of Galt, which uses UDP-glucose for galactose catabolism, was higher only in male mice, whereas Ugdh, which uses UDP-glucose for glucuronate biosynthesis, was higher only in female mice (Figure 7E).

Targeted metabolomics of GLUT3 KD neurons at 1 and 24 h post incubation with [U-¹³C]glucose revealed increased amounts of UDP-glucose and glucose/fructose-6P (G6P-F6P) after 1 h (Figure 7F). As the metabolites in the media were consumed over the 24-h period, the total amount of UDP-glucose and G6P-F6P in GLUT3 KD neurons declined. The change in total UDP-glucose amount between the 1- and 24-h time points indicates a higher rate of UDP-glucose consumption relative to its biosynthesis; the increased expression of Gale could reflect a response to meet the catabolic demand for UDP-glucose through its increased production. Given that the B27 supplement used in the neuronal differentiation media contains galactose (0.04 mM in media; Figure S7E), we investigated whether human neurons can metabolize galactose via glycolysis and the TCA cycle. Incubation with [U-¹³C]galactose resulted in robust ¹³C-incorporation by both NTG and GLUT3 KD neurons into galactose-1P, glucose-6P, as well as into glycolytic, PPP, and TCA metabolites (Figures 7G–7J and S7F). Indicative of up-regulated galactose metabolism in response to GLUT3 KD, total levels of galactose-1P and glucose-6P in GLUT3 KD neurons (\approx 80% KD by rtPCR) cultured with [U-¹³C]galactose were higher than in NTG neurons (Figure 7H). GLUT3 KD neurons also had slightly higher fractions of ¹³C-labeling in alpha-ketoglutarate (α -KG) and malate, relative to controls (Figure S7F).

To further assess the interplay between glucose and galactose, we examined the impact of equimolar amounts of galactose and glucose on glucose metabolism. Addition of galactose increased the total levels of galactose-1P and glucose-6P, while markedly decreasing the percentage labeling of galactose-1P and glucose-6P following a 24-h incubation with [U-¹³C]glucose (Figure 7K), consistent with galactose competing with glucose for conversion into these metabolites. Moreover, the addition of galactose contributed to the increased levels of PPP metabolites 6P-gluconate and ribose-5P (Figure 7L). Interestingly, while the addition of galactose decreased the fractional labeling from [U-¹³C]glucose in upstream glycolytic metabolites 3PG and PEP, labeling in both pyruvate and lactate remained unchanged (Figure 7M). However, the total amounts of pyruvate and lactate increased \sim 60% with the addition of galactose only in the GLUT3 KD neurons. The total amounts of TCA metabolites citrate, α -KG, and malate increased in both NTG and GLUT3 KD neurons with the addition of galactose despite the fractional labeling from glucose remaining the same (Figure S7G). The contrast in fractional [U-¹³C]glucose labeling between the PPP and glycolysis with the addition of galactose illustrates that galactose is preferentially shunted through the PPP, while glucose is directed through glycolysis to the TCA cycle. Taken together, the differentially expressed metabolic and mitochondrial genes suggest a multi-pronged compensatory response

to GLUT3cKO involving a shift toward mitochondrial respiration and enhanced galactose metabolism.

DISCUSSION

It has been proposed that neurons derive most or all of their ATP from aerobic respiration,⁶⁸ and the extent to which neurons directly take up and metabolize glucose through glycolysis, and whether this is required, especially under physiologic conditions, is unknown. Here we show that neurons can indeed take up and metabolize glucose through glycolysis and use this to produce the majority of their TCA metabolites. Moreover, we show they require glucose uptake to maintain ATP levels at the synapse, and normal neuronal function *in vivo*. Our findings also reveal systems-level sex differences in glucose metabolism and identify increased galactose metabolism as a mechanism to compensate for impaired neuronal glucose uptake.

Neurons require direct glucose uptake and metabolism through glycolysis

Multiple lines of evidence suggest that neurons directly import glucose in culture.^{6,52,53,69} However, direct confirmation of this has remained elusive because of the presence of glia in primary rodent neuron cultures, which limits the ability to distinguish glial from neuronal uptake. Here, we used metabolomics on near homogeneous human neuron cultures to demonstrate that neurons do indeed directly import glucose. Moreover, we show that GLUT3 is required for normal glucose uptake, but the effect is most robust when glucose levels are at physiologic brain levels. At the higher glucose levels typically used for neuronal culture, glucose could still enter GLUT3KO neurons and reach WT levels. The mechanism may involve other GLUTs, although we found no evidence of upregulation of other GLUTs in GLUT3cKO mice.

What are the consequences of decreased glucose import into neurons? We found that GLUT3KO neurons are less metabolically responsive to changes in extracellular glucose, and that decreased glucose import in GLUT3KO neurons results in a marked decrease in glucose levels during neural activity. Our metabolomics data also suggest that, when glucose uptake is limited, neurons may prioritize maintaining flux of glucose metabolites through the TCA cycle at the expense of both aerobic glycolysis (lactate) and the PPP.

Ultimately, decreased glucose uptake into neurons impairs function, as mice with postnatal deletion of GLUT3 develop spatial learning and memory deficits. Our findings in GLUT3cKO mice are consistent with and build on findings from Shin et al., who observed memory deficits in male mice in a similar model of GLUT3 deficiency.⁷ These deficits may result from synaptic dysfunction, as we found that GLUT3 deficiency compromised ATP levels at the nerve terminal but not the cell body. Notably, cell body and dendrite function may also be compromised *in vivo*, especially as decreased glucose levels in GLUT3KO neurons were only apparent at low glucose levels. However, we did not detect deficits *ex vivo* under higher (10 mM) glucose conditions.

Another important question is how many of the above changes can be attributed to deficits in glycolysis versus other glucose-supported functions, such as the PPP. Even if glucose is taken

up by neurons, neurons have been hypothesized to metabolize glucose primarily through the PPP due to insufficient phosphofructokinase to support glycolysis.^{8,9} Moreover, neuronal glycolysis is reported to increase in response to certain pathologic stressors, and pharmacologically inhibiting this increase may be neuroprotective.^{10–12} Here, we show that human neurons do indeed metabolize glucose through glycolysis and can derive most of their TCA metabolites from glucose. Moreover, we show that postnatal deletion of PKM1 in neurons *in vivo* resulted in memory loss from 12 months of age in female mice, while males were unaffected at this age. This was accompanied by increased conversion of HP [1-¹³C]pyruvate to [1-¹³C]lactate, indicating a functionally significant disruption in glycolytic metabolism upon PKM1 deletion, compelling evidence that neurons (or at least neurons from females) require glycolysis for normal function.

Importantly, our findings do not exclude the lactate shuttle as an important contributor to the bioenergetic function of neurons. Indeed, considerable evidence shows both that neurons can support neural activity with either respiration or glycolysis, and that neurons and other cell types can readily switch between fuel sources depending on their energy requirements and local substrate availability.^{6,23,52,53} However, our data indicate that the lactate shuttle is unable to upregulate and fully support neuronal energy requirements on its own.

Sex-specific requirements for glucose uptake and glycolysis

Interestingly, the PKM1 metabolic deficits detected by HP ¹³C MR occurred exclusively in females, consistent with the memory deficit specific to female PKM1cKO mice. Interpretation of this sex-specific HP change is complex, but likely reflects not only the primary disruption in neuronal glycolysis but also potential compensatory changes in glial metabolism and neuro-glial coupling (lactate shuttle). The HP pyruvate and lactate signals encompass vascular, neuronal, and glial components, and the MR acquisition parameters (in particular flip angles⁷⁰) are likely to play a crucial role in the relative contribution of each to the HP lactate-to-pyruvate ratios measured by MR. Although future studies are needed to fully understand the mechanisms driving the HP changes, this represents (to our knowledge) only the second report of HP metabolic imaging detecting sex differences *in vivo*,⁶³ and the first one in the brain. Our data support the potential of HP metabolic imaging to reveal changes in brain energy metabolism with aging and disease.⁷¹ Notably, [¹⁸F]FDG-PET imaging did not detect any changes following either GLUT3 or PKM1 deletion, which might be due to high background levels, masking changes occurring in the relatively small CA1 area, and might also indicate that most [¹⁸F]FDG-PET signal originates from glial cells.⁷²

We hypothesize that PKM1cKO males may also eventually develop motor deficits, and point out that the human neurons with PKM KD (both PKM1 and M2 isoforms), in which we observed metabolic differences, are male in origin. Nonetheless, neurons in female mice were more susceptible to PKM1 inhibition in our paradigm, perhaps suggesting sex-specific differences in compensatory mechanisms. These differences may also apply to other tissues, as loss of PKM2 promotes hepatocellular cancer at a far greater rate in male than female mice.⁷³

GLUT3cKO mice also developed broad sex-specific changes. First, GLUT3cKO female mice had decreased body weight, consistent with a prior finding that GLUT3 deletion in neurons and glia during early embryogenesis decreased body weight in females but not males.⁷⁴ Remarkably, our study replicates these results when GLUT3 is deleted in neuronal populations not normally associated with weight regulation. Although we have not excluded the possibility that changes in body weight resulted from memory deficits leading to decreased food consumption, the memory changes occurred in both male and female mice, while the changes in body weight and brain volume were only in females. Moreover, the memory changes involved only the GLUT3cKO mice, while the weight changes also occurred in heterozygotes. An independent study showed that mice with embryonic deletion of GLUT3 throughout the brain had decreased body weight at postnatal day 15 (males and females were analyzed together), despite increased food intake,⁷ supporting that the changes in body weight in our model are unlikely to result from insufficient food intake. Interestingly, hippocampal sharp wave-ripples were recently found to regulate peripheral glucose concentrations via a pathway projecting to the lateral septum and hypothalamus,³⁶ raising the possibility that GLUT3cKO might influence food consumption through alternate mechanisms. However, the role of this pathway in our observations is unclear, and peripheral glucose levels were unchanged in GLUT3cKO mice.

Metabolic rewiring in response to glucose deficiency

A critical question is how neurons compensate when glucose is limiting *in vivo*. Indeed, although the capacity for metabolic rewiring has been considered limited in neurons,⁷⁵ broad compensatory mechanisms to maintain TCA metabolites are observed in the setting of mitochondrial dysfunction.⁷⁶ Here, we identify two potential mechanisms of neuronal compensation. First, we show that GLUT3cKO neurons upregulate several mitochondrial genes that can promote mitochondrial respiration, including *Mitf*, *lars2*, and *Echdc2*. Upregulation of these genes may reflect an attempt to boost mitochondrial respiration by increasing the utilization of available glucose and by boosting respiration generated from other fuel sources.

Interestingly, another gene that may promote mitochondrial respiration, *PDP2*, was increased in male (but not female) GLUT3cKO neurons, perhaps contributing to the increased susceptibility of female mice to GLUT3 loss. Indeed, considerable evidence indicates that males and females (mice and humans) have fundamental differences in glucose and energy metabolism,⁷⁷ and neurons and skeletal muscle from females rely more on fatty acid β -oxidation as an energy source than those from males.^{78,79} However, the molecular basis for these differences is poorly understood. Another gene that responded differently between sexes to GLUT3cKO was *Bcs1l*, a mitochondrial protein responsible for the final assembly of complex III; mutations to *Bcs1l* have been causally linked to complex III deficiency,⁸⁰ impaired β -oxidation,⁸¹ and hypoglycemia.⁸² Additional research is required to determine whether the differential expression of such genes may underlie some of the differences in energy metabolism between sexes.

In addition to genes that regulate respiration, we also identified changes in genes regulating galactose metabolism. Galactose catabolism via the Leloir pathway yields glucose 6-phosphate to feed into both glycolysis and the PPP.⁶⁶ We show that boosting galactose metabolism is an alternate mechanism to bypass glucose deficiency. Notably, GLUT3 not only mediates glucose transport, it also transports other carbohydrates, including galactose.⁸³ As such, upregulation of galactose metabolism genes in the setting of GLUT3 loss of function may, in part, be a compensatory response to maintain intraneuronal galactose levels. However, our finding that galactose import is upregulated in response to GLUT3KD indicates that galactose can enter neurons through other mechanisms, and shows that neurons do indeed upregulate galactose metabolism to meet energy demands when glucose is insufficient. Indeed, GLUT3 has an extremely low affinity for galactose, and galactose is a substrate for other GLUTs (GLUT1, GLUT8) that hippocampal neurons express at lower levels.^{84,85} As such, our data provide some of the first evidence that neurons normally metabolize galactose, suggesting that galactose may contribute to normal brain energy homeostasis and may be able to compensate when glucose metabolism is disrupted. However, given the changes in memory, brain size, and body weight in GLUT3cKO mice, it is clear that this mechanism was not sufficient to compensate for decreased glucose uptake in CA1 neurons, at least with physiological levels of energy substrates.

Implications for neurologic and neurodegenerative diseases

Our findings also suggest that impairments in neuronal glucose metabolism may directly contribute to neuronal dysfunction and degeneration in a range of conditions of insufficient brain glucose, such as hypoglycemia, stroke, and Alzheimer's disease (AD). In AD, changes in glucose metabolism could result from a primary disruption of glucose uptake or glycolysis, or may reflect secondary compensatory changes to other metabolic deficits or to changes in neural activity that affect energy requirements. Interestingly, brains of patients with AD have markedly decreased levels of GLUT3 in the hippocampus and cortex even when normalized for synaptic loss,⁸⁶ raising the possibility that decreased GLUT3 contributes to the decreased glucose uptake in AD. Moreover, overexpression of human GLUT3 is protective in a model of amyotrophic lateral sclerosis.⁸⁷ As such, the findings presented here provide important insights into the basic biology of neuronal glucose metabolism and how changes in glucose metabolism may contribute to neurologic disease.

Limitations of the study

The metabolomics experiments prove that human neurons metabolize glucose through glycolysis. However, the pure neuronal population studied lacks glial cells, circuits, and three-dimensional interactions found *in vivo* that likely influence metabolism. Therefore, the relative flux of glucose metabolites through glycolysis versus the PPP, and through downstream glycolytic pathways, may differ between the human neurons studied and human and mouse neurons *in vivo*.

Although our paper focuses on changes in neurons, the HP pyruvate and MRI studies *in vivo* lack cell-type resolution. We have

begun to address this by postnatally deleting GLUT3 and PKM1 specifically in neurons. These studies show that neurons need glycolysis to maintain normal brain energy metabolism and volume. However, they do not definitively prove that the changes in either HP pyruvate metabolism or brain volume directly result from changes in the neurons themselves rather than via non-cell-autonomous effects on glia. Moreover, the requirement for neuronal glycolysis does not provide insight into whether neurons also require astrocyte-derived lactate and, if so, the balance of energy derived from neuronal glucose metabolism versus astrocyte-derived lactate.

Last, our work begins to address the key question of how neurons compensate when glucose is limiting. We identify a compensatory upregulation of galactose metabolism that fuels downstream energy metabolites as well as several changes in the expression of metabolism-based genes, but additional studies are required to determine which of these changes are most critical in restoring energy.

STAR★METHODS

Detailed methods are provided in the online version of this paper and include the following:

- KEY RESOURCES TABLE
- RESOURCE AVAILABILITY
 - Lead contact
 - Materials availability
 - Data and code availability
- EXPERIMENTAL MODEL AND SUBJECT DETAILS
 - Mice
 - Generation of iPSC lines and iPSC-derived neuronal cultures
 - Primary neuronal cultures
- METHOD DETAILS
 - Metabolomics
 - Galactose media measurements
 - Behavioral testing
 - Histology and immunostaining
 - *In vivo* MR acquisitions
 - *Ex vivo* MRI acquisitions
 - MR data analysis
 - *In vivo* PET acquisitions
 - PET data analysis
 - Electrophysiology
 - Live imaging of glucose or ATP
 - Blood glucose level measurement
 - Spatial gene expression
- QUANTIFICATION AND STATISTICAL ANALYSIS

SUPPLEMENTAL INFORMATION

Supplemental information can be found online at <https://doi.org/10.1016/j.celrep.2023.112335>.

ACKNOWLEDGMENTS

We thank Jonathan Marvin (HHMI, Janelia) for providing the glucose sensor, Jessica Speckart and Julia Holtzman and the Gladstone Behavioral core for

assistance with behavioral testing and data analysis, and Michael Gil for comments on the manuscript. We also thank Johanna ten Hoeve, Thomas G. Graeber, and the UCLA Metabolomics Center for their assistance with metabolomics studies and data processing. We also thank Blaise Ndjamen and Anna Torrent Moreno of the Gladstone Histology and Light Microscopy core, Eric Chow and the UCSF Center for Advanced Technology, Mylinh Bernardi and the Gladstone Genomics Core, as well as Kathryn Claiborn for helping edit the manuscript and Erica Delin for administrative assistance. This work was supported by National Institutes of Health RF1 AG064170 (to K.N. and M.M.C.), R01 AG065428 (K.N.), AG065428-03S1 (K.N., M.K.), R01 NS102156 (M.M.C.), R21 AI153749 (M.M.C., C.G.), National Institute on Aging R01 AG061150 (M.Z.), R01 AG071697 and P01 AG073082 (Y.H.), R01 CA168653 and R35 CA242379 (M.G.V.H.), R01 DK105550 (J.C.R.), K01AG078485 (N.B.), and National Institutes of Health RR18928 (Gladstone Institutes). It was also supported by the UCSF Bakar Aging Research Institute (BARI) (K.N.), the Alzheimer's Association (H.L. and L.M.L.P.), a BrightFocus Foundation Award (L.M.L.P.), a Berkelhammer Award for Excellence in Neuroscience (Y.J.S.), Hillblom Center and Bakar Aging Research Institute Graduate Fellowship (J.Y.), and a Chan Zuckerberg Initiative Neurodegeneration Challenge Network Ben Barres Early Career Acceleration Award (M.K.).

AUTHOR CONTRIBUTIONS

H.L., C.G., Y.J.S., M.Z., M.M.C., and K. Nakamura designed the research. H.L., C.G., Y.J.S., M.Z., L.M.L.P., L.S., J.Y., K. Nguyen, B.T., X.G., N.B., and I.L. performed research. T.L.D., M.K., Y.H., J.C.R., and M.V.H. provided guidance and reagents for critical techniques. H.L., C.G., Y.J.S., M.Z., L.M.L.P., J.Y., K. Nguyen, I.L., M.M.C., and K. Nakamura analyzed data. H.L., C.G., Y.J.S., M.M.C., and K. Nakamura wrote the paper with assistance from all co-authors.

DECLARATION OF INTERESTS

M.G.V.H. discloses that he is a scientific advisor for Agios Pharmaceuticals, iTeos Therapeutics, Sage Therapeutics, Faeth Therapeutics, DRIOA Ventures, and Auron Therapeutics. Y.H. discloses that he is a co-founder and scientific advisory board member of GABAeron.

INCLUSION AND DIVERSITY

We support inclusive, diverse, and equitable conduct of research.

Received: January 3, 2023

Revised: February 22, 2023

Accepted: March 20, 2023

Published: April 6, 2023

REFERENCES

- Bélanger, M., Allaman, I., and Magistretti, P.J. (2011). Brain energy metabolism: focus on astrocyte-neuron metabolic cooperation. *Cell Metab.* *14*, 724–738. <https://doi.org/10.1016/j.cmet.2011.08.016>.
- Patel, A.B., Lai, J.C.K., Chowdhury, G.M.I., Hyder, F., Rothman, D.L., Shulman, R.G., and Behar, K.L. (2014). Direct evidence for activity-dependent glucose phosphorylation in neurons with implications for the astrocyte-to-neuron lactate shuttle. *Proc. Natl. Acad. Sci. USA* *111*, 5385–5390. <https://doi.org/10.1073/pnas.1403576111>.
- Lundgaard, I., Li, B., Xie, L., Kang, H., Sanggaard, S., Haswell, J.D.R., Sun, W., Goldman, S., Blekot, S., Nielsen, M., et al. (2015). Direct neuronal glucose uptake heralds activity-dependent increases in cerebral metabolism. *Nat. Commun.* *6*, 6807. <https://doi.org/10.1038/ncomms7807>.
- Ivanov, A.I., Malkov, A.E., Waseem, T., Mukhtarov, M., Buldakova, S., Gubkina, O., Zilberter, M., and Zilberter, Y. (2014). Glycolysis and oxidative phosphorylation in neurons and astrocytes during network activity in hippocampal slices. *J. Cereb. Blood Flow Metab.* *34*, 397–407. <https://doi.org/10.1038/jcbfm.2013.222>.
- Maier, F. (1995). Immunolocalization of GLUT1 and GLUT3 glucose transporters in primary cultured neurons and glia. *J. Neurosci. Res.* *42*, 459–469. <https://doi.org/10.1002/jnr.490420404>.
- Ashrafi, G., Wu, Z., Farrell, R.J., and Ryan, T.A. (2017). GLUT4 mobilization supports energetic demands of active synapses. *Neuron* *93*, 606–615.e3. <https://doi.org/10.1016/j.neuron.2016.12.020>.
- Shin, B.C., Cepeda, C., Estrada-Sánchez, A.M., Levine, M.S., Hodaei, L., Dai, Y., Jung, J., Ganguly, A., Clark, P., and Devaskar, S.U. (2018). Neural deletion of glucose transporter isoform 3 creates distinct postnatal and adult neurobehavioral phenotypes. *J. Neurosci.* *38*, 9579–9599. <https://doi.org/10.1523/JNEUROSCI.0503-18.2018>.
- Herrero-Mendez, A., Almeida, A., Fernández, E., Maestre, C., Moncada, S., and Bolaños, J.P. (2009). The bioenergetic and antioxidant status of neurons is controlled by continuous degradation of a key glycolytic enzyme by APC/C-Cdh1. *Nat. Cell Biol.* *11*, 747–752. <https://doi.org/10.1038/ncb1881>.
- Bolaños, J.P., Almeida, A., and Moncada, S. (2010). Glycolysis: a bioenergetic or a survival pathway? *Trends Biochem. Sci.* *35*, 145–149. <https://doi.org/10.1016/j.tibs.2009.10.006>.
- Rodríguez-Rodríguez, P., Fernández, E., Almeida, A., and Bolaños, J.P. (2012). Excitotoxic stimulus stabilizes PFKFB3 causing pentose-phosphate pathway to glycolysis switch and neurodegeneration. *Cell Death Differ.* *19*, 1582–1589. <https://doi.org/10.1038/cdd.2012.33>.
- Burmistrova, O., Olias-Arjona, A., Lapresa, R., Jimenez-Blasco, D., Ereemeeva, T., Shishov, D., Romanov, S., Zakurdaeva, K., Almeida, A., Fedichev, P.O., et al. (2019). Targeting PFKFB3 alleviates cerebral ischemia-reperfusion injury in mice. *Sci. Rep.* *9*, 11670. <https://doi.org/10.1038/s41598-019-48196-z>.
- Lopez-Fabuel, I., Garcia-Macia, M., Buondelmonte, C., Burmistrova, O., Bonora, N., Alonso-Batan, P., Morant-Ferrando, B., Vicente-Gutierrez, C., Jimenez-Blasco, D., Quintana-Cabrera, R., et al. (2022). Aberrant up-regulation of the glycolytic enzyme PFKFB3 in CLN7 neuronal ceroid lipofuscinosis. *Nat. Commun.* *13*, 536. <https://doi.org/10.1038/s41467-022-28191-1>.
- Díaz-García, C.M., Mongeon, R., Lahmann, C., Koveal, D., Zucker, H., and Yellen, G. (2017). Neuronal stimulation triggers neuronal glycolysis and not lactate uptake. *Cell Metab.* *26*, 361–374.e4. <https://doi.org/10.1016/j.cmet.2017.06.021>.
- Ardenjaer-Larsen, J.H., Fridlund, B., Gram, A., Hansson, G., Hansson, L., Lerche, M.H., Servin, R., Thaning, M., and Golman, K. (2003). Increase in signal-to-noise ratio of > 10,000 times in liquid-state NMR. *Proc. Natl. Acad. Sci. USA* *100*, 10158–10163. <https://doi.org/10.1073/pnas.1733835100>.
- Brindle, K.M., Bohndiek, S.E., Gallagher, F.A., and Kettunen, M.I. (2011). Tumor imaging using hyperpolarized ¹³C magnetic resonance spectroscopy. *Magn. Reson. Med.* *66*, 505–519. <https://doi.org/10.1002/mrm.22999>.
- Chaumeil, M.M., Najac, C., and Ronen, S.M. (2015). Studies of metabolism using (¹³C) MRS of hyperpolarized probes. *Methods Enzymol.* *567*, 1–71. <https://doi.org/10.1016/bs.mie.2015.04.001>.
- Nelson, S.J., Kurhanewicz, J., Vigneron, D.B., Larson, P.E.Z., Harzstark, A.L., Ferrone, M., van Criekinge, M., Chang, J.W., Bok, R., Park, I., et al. (2013). Metabolic imaging of patients with prostate cancer using hyperpolarized [1-(¹³C)]pyruvate. *Sci. Transl. Med.* *5*, 198ra108. <https://doi.org/10.1126/scitranslmed.3006070>.
- Gallagher, F.A., Woitek, R., McLean, M.A., Gill, A.B., Manzano Garcia, R., Provenzano, E., Riemer, F., Kaggie, J., Chhabra, A., Ursprung, S., et al. (2020). Imaging breast cancer using hyperpolarized carbon-13 MRI. *Proc. Natl. Acad. Sci. USA* *117*, 2092–2098. <https://doi.org/10.1073/pnas.1913841117>.
- Park, I., Larson, P.E.Z., Gordon, J.W., Carvajal, L., Chen, H.Y., Bok, R., Van Criekinge, M., Ferrone, M., Slater, J.B., Xu, D., et al. (2018). Development of methods and feasibility of using hyperpolarized carbon-13

- imaging data for evaluating brain metabolism in patient studies. *Magn. Reson. Med.* 80, 864–873. <https://doi.org/10.1002/mrm.27077>.
20. Wang, C., Ward, M.E., Chen, R., Liu, K., Tracy, T.E., Chen, X., Xie, M., Sohn, P.D., Ludwig, C., Meyer-Franke, A., et al. (2017). Scalable production of iPSC-derived human neurons to identify tau-lowering compounds by high-content screening. *Stem Cell Rep.* 9, 1221–1233. <https://doi.org/10.1016/j.stemcr.2017.08.019>.
 21. Zhang, Y., Pak, C., Han, Y., Ahlenius, H., Zhang, Z., Chanda, S., Marro, S., Patzke, C., Acuna, C., Covy, J., et al. (2013). Rapid single-step induction of functional neurons from human pluripotent stem cells. *Neuron* 78, 785–798. <https://doi.org/10.1016/j.neuron.2013.05.029>.
 22. Tian, R., Gachechiladze, M.A., Ludwig, C.H., Laurie, M.T., Hong, J.Y., Nathaniel, D., Prabhu, A.V., Fernandopulle, M.S., Patel, R., Abshari, M., et al. (2019). CRISPR interference-based platform for multimodal genetic screens in human iPSC-derived neurons. *Neuron* 104, 239–255.e12. <https://doi.org/10.1016/j.neuron.2019.07.014>.
 23. Bennett, N.K., Nguyen, M.K., Darch, M.A., Nakaoka, H.J., Cousineau, D., Ten Hoeve, J., Graeber, T.G., Schuelke, M., Maltepe, E., Kampmann, M., et al. (2020). Defining the ATPome reveals cross-optimization of metabolic pathways. *Nat. Commun.* 11, 4319. <https://doi.org/10.1038/s41467-020-18084-6>.
 24. Contat, C., Ancy, P.B., Zangger, N., Sabatino, S., Pascual, J., Escrig, S., Jensen, L., Goepfert, C., Lanz, B., Lepore, M., et al. (2020). Combined deletion of Glut1 and Glut3 impairs lung adenocarcinoma growth. *Elife* 9, e53618. <https://doi.org/10.7554/eLife.53618>.
 25. Horn, K.E., Glasgow, S.D., Gobert, D., Bull, S.J., Luk, T., Girgis, J., Tremblay, M.E., McEachern, D., Bouchard, J.F., Haber, M., et al. (2013). DCC expression by neurons regulates synaptic plasticity in the adult brain. *Cell Rep.* 3, 173–185. <https://doi.org/10.1016/j.celrep.2012.12.005>.
 26. Tsien, J.Z., Chen, D.F., Gerber, D., Tom, C., Mercer, E.H., Anderson, D.J., Mayford, M., Kandel, E.R., and Tonegawa, S. (1996). Subregion- and cell type-restricted gene knockout in mouse brain. *Cell* 87, 1317–1326.
 27. Bergsman, J.B., and Tsien, R.W. (2000). Syntaxin modulation of calcium channels in cortical synaptosomes as revealed by botulinum toxin C1. *J. Neurosci.* 20, 4368–4378.
 28. Miller, J.A., Guillozet-Bongaarts, A., Gibbons, L.E., Postupna, N., Renz, A., Beller, A.E., Sunkin, S.M., Ng, L., Rose, S.E., Smith, K.A., et al. (2017). Neuropathological and transcriptomic characteristics of the aged brain. *Elife* 6, e31126. <https://doi.org/10.7554/eLife.31126>.
 29. Lesburguères, E., Sparks, F.T., O'Reilly, K.C., and Fenton, A.A. (2016). Active place avoidance is no more stressful than unreinforced exploration of a familiar environment. *Hippocampus* 26, 1481–1485. <https://doi.org/10.1002/hipo.22666>.
 30. Shields, L.Y., Kim, H., Zhu, L., Haddad, D., Berthet, A., Pathak, D., Lam, M., Ponnusamy, R., Diaz-Ramirez, L.G., Gill, T.M., et al. (2015). Dynamically related protein 1 is required for normal mitochondrial bioenergetic and synaptic function in CA1 hippocampal neurons. *Cell Death Dis.* 6, e1725. <https://doi.org/10.1038/cddis.2015.94>.
 31. Marty, S., and Peschanski, M. (1995). Effects of target deprivation on the morphology and survival of adult dorsal column nuclei neurons. *J. Comp. Neurol.* 356, 523–536. <https://doi.org/10.1002/cne.903560404>.
 32. Li, Z., Okamoto, K.I., Hayashi, Y., and Sheng, M. (2004). The importance of dendritic mitochondria in the morphogenesis and plasticity of spines and synapses. *Cell* 119, 873–887.
 33. Tonegawa, S., Tsien, J.Z., McHugh, T.J., Huerta, P., Blum, K.I., and Wilson, M.A. (1996). Hippocampal CA1-region-restricted knockout of NMDAR1 gene disrupts synaptic plasticity, place fields, and spatial learning. *Cold Spring Harb. Symp. Quant. Biol.* 61, 225–238.
 34. Wang, X., Zhang, C., Szábo, G., and Sun, Q.Q. (2013). Distribution of CaMKIIalpha expression in the brain in vivo, studied by CaMKIIalpha-GFP mice. *Brain Res.* 1518, 9–25. <https://doi.org/10.1016/j.brainres.2013.04.042>.
 35. Semple, B.D., Blomgren, K., Gimlin, K., Ferriero, D.M., and Noble-Haueslein, L.J. (2013). Brain development in rodents and humans: identifying benchmarks of maturation and vulnerability to injury across species. *Prog. Neurobiol.* 106–107, 1–16. <https://doi.org/10.1016/j.pneurobio.2013.04.001>.
 36. Tingley, D., McClain, K., Kaya, E., Carpenter, J., and Buzsáki, G. (2021). A metabolic function of the hippocampal sharp wave-ripple. *Nature* 597, 82–86. <https://doi.org/10.1038/s41586-021-03811-w>.
 37. Macdonald, E.B., Begovatz, P., Barton, G.P., Erickson-Bhatt, S., Inman, D.R., Cox, B.L., Eliceiri, K.W., Strigel, R.M., Ponik, S.M., and Fain, S.B. (2021). Hyperpolarized ¹³C magnetic resonance spectroscopic imaging of pyruvate metabolism in murine breast cancer models of different metastatic potential. *Metabolites* 11, 274. <https://doi.org/10.3390/metabo11050274>.
 38. Perkons, N.R., Kiefer, R.M., Noji, M.C., Pourfathi, M., Ackerman, D., Siddiqui, S., Tischfield, D., Profka, E., Johnson, O., Pickup, S., et al. (2020). Hyperpolarized metabolic imaging detects latent hepatocellular carcinoma domains surviving locoregional therapy. *Hepatology* 72, 140–154. <https://doi.org/10.1002/hep.30970>.
 39. Varma, G., Seth, P., Coutinho de Souza, P., Callahan, C., Pinto, J., Vaidya, M., Sonzogni, O., Sukhatme, V., Wulf, G.M., and Grant, A.K. (2021). Visualizing the effects of lactate dehydrogenase (LDH) inhibition and LDH-A genetic ablation in breast and lung cancer with hyperpolarized pyruvate NMR. *NMR Biomed.* 34, e4560. <https://doi.org/10.1002/nbm.4560>.
 40. Guglielmetti, C., Najac, C., Didonna, A., Van der Linden, A., Ronen, S.M., and Chaumeil, M.M. (2017). Hyperpolarized ¹³C MR metabolic imaging can detect neuroinflammation in vivo in a multiple sclerosis murine model. *Proc. Natl. Acad. Sci. USA* 114, E6982–E6991. <https://doi.org/10.1073/pnas.1613345114>.
 41. Harris, R.A., Lone, A., Lim, H., Martinez, F., Frame, A.K., Scholl, T.J., and Cumming, R.C. (2019). Aerobic glycolysis is required for spatial memory acquisition but not memory retrieval in mice. *eNeuro* 6, ENEURO.0389-18.2019. <https://doi.org/10.1523/ENEURO.0389-18.2019>.
 42. Rao, Y., Gammon, S., Zacharias, N.M., Liu, T., Salzillo, T., Xi, Y., Wang, J., Bhattacharya, P., and Piwnicka-Worms, D. (2020). Hyperpolarized [1-(¹³C)] pyruvate-to-[1-(¹³C)] lactate conversion is rate-limited by monocarboxylate transporter-1 in the plasma membrane. *Proc. Natl. Acad. Sci. USA* 117, 22378–22389. <https://doi.org/10.1073/pnas.2003537117>.
 43. Day, S.E., Kettunen, M.I., Gallagher, F.A., Hu, D.E., Lerche, M., Wolber, J., Golman, K., Ardenkjaer-Larsen, J.H., and Brindle, K.M. (2007). Detecting tumor response to treatment using hyperpolarized ¹³C magnetic resonance imaging and spectroscopy. *Nat. Med.* 13, 1382–1387. <https://doi.org/10.1038/nm1650>.
 44. Brindle, K.M. (2015). Imaging metabolism with hyperpolarized ¹³C-labeled cell substrates. *J. Am. Chem. Soc.* 137, 6418–6427. <https://doi.org/10.1021/jacs.5b03300>.
 45. Witney, T.H., Kettunen, M.I., Day, S.E., Hu, D.E., Neves, A.A., Gallagher, F.A., Fulton, S.M., and Brindle, K.M. (2009). A comparison between radio-labeled fluorodeoxyglucose uptake and hyperpolarized ¹³C-labeled pyruvate utilization as methods for detecting tumor response to treatment. *Neoplasia* 11, 574–582. <https://doi.org/10.1593/neo.09254>.
 46. Xie, N., Zhang, L., Gao, W., Huang, C., Huber, P.E., Zhou, X., Li, C., Shen, G., and Zou, B. (2020). NAD(+) metabolism: pathophysiologic mechanisms and therapeutic potential. *Signal Transduct. Target. Ther.* 5, 227. <https://doi.org/10.1038/s41392-020-00311-7>.
 47. Davidson, S.M., Schmidt, D.R., Heyman, J.E., O'Brien, J.P., Liu, A.C., Israelsen, W.J., Dayton, T.L., Sehgal, R., Bronson, R.T., Freinkman, E., et al. (2022). Pyruvate kinase M1 suppresses development and progression of prostate adenocarcinoma. *Cancer Res.* 82, 2403–2416. <https://doi.org/10.1158/0008-5472.CAN-21-2352>.
 48. Zhang, Y., Chen, K., Sloan, S.A., Bennett, M.L., Scholze, A.R., O'Keefe, S., Phatnani, H.P., Guarnieri, P., Caneda, C., Ruderisch, N., et al. (2014). An RNA-sequencing transcriptome and splicing database of glia, neurons, and vascular cells of the cerebral cortex. *J. Neurosci.* 34, 11929–11947. <https://doi.org/10.1523/JNEUROSCI.1860-14.2014>.

49. Tech, K., Tikunov, A.P., Farooq, H., Morrissy, A.S., Meidinger, J., Fish, T., Green, S.C., Liu, H., Li, Y., Mungall, A.J., et al. (2017). Pyruvate kinase inhibits proliferation during postnatal cerebellar neurogenesis and suppresses medulloblastoma formation. *Cancer Res.* 77, 3217–3230. <https://doi.org/10.1158/0008-5472.CAN-16-3304>.
50. Park, I., Mukherjee, J., Ito, M., Chaumeil, M.M., Jalbert, L.E., Gaensler, K., Ronen, S.M., Nelson, S.J., and Pieper, R.O. (2014). Changes in pyruvate metabolism detected by magnetic resonance imaging are linked to DNA damage and serve as a sensor of temozolomide response in glioblastoma cells. *Cancer Res.* 74, 7115–7124. <https://doi.org/10.1158/0008-5472.CAN-14-0849>.
51. Casson, R.J., Wood, J.P.M., Han, G., Kittipassorn, T., Peet, D.J., and Chidlow, G. (2016). M-Type pyruvate kinase isoforms and lactate dehydrogenase A in the mammalian retina: metabolic implications. *Invest. Ophthalmol. Vis. Sci.* 57, 66–80. <https://doi.org/10.1167/iovs.15-17962>.
52. Pathak, D., Shields, L.Y., Mendelsohn, B.A., Haddad, D., Lin, W., Gerencser, A.A., Kim, H., Brand, M.D., Edwards, R.H., and Nakamura, K. (2015). The role of mitochondrially derived ATP in synaptic vesicle recycling. *J. Biol. Chem.* 290, 22325–22336. <https://doi.org/10.1074/jbc.M115.656405>.
53. Rangaraju, V., Calloway, N., and Ryan, T.A. (2014). Activity-driven local ATP synthesis is required for synaptic function. *Cell* 156, 825–835. <https://doi.org/10.1016/j.cell.2013.12.042>.
54. Vannucci, S.J., Maher, F., and Simpson, I.A. (1997). Glucose transporter proteins in brain: delivery of glucose to neurons and glia. *Glia* 21, 2–21. [https://doi.org/10.1002/\(sici\)1098-1136](https://doi.org/10.1002/(sici)1098-1136).
55. Imamura, H., Nhat, K.P.H., Togawa, H., Saito, K., Iino, R., Kato-Yamada, Y., Nagai, T., and Noji, H. (2009). Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators. *Proc. Natl. Acad. Sci. USA* 106, 15651–15656. <https://doi.org/10.1073/pnas.0904764106>.
56. Mendelsohn, B.A., Bennett, N.K., Darch, M.A., Yu, K., Nguyen, M.K., Puciarelli, D., Nelson, M., Horlbeck, M.A., Gilbert, L.A., Hyun, W., et al. (2018). A high-throughput screen of real-time ATP levels in individual cells reveals mechanisms of energy failure. *PLoS Biol.* 16, e2004624. <https://doi.org/10.1371/journal.pbio.2004624>.
57. Vona, B., Maroofian, R., Bellacchio, E., Najafi, M., Thompson, K., Alahmad, A., He, L., Ahangari, N., Rad, A., Shahrokhzadeh, S., et al. (2018). Expanding the clinical phenotype of IARS2-related mitochondrial disease. *BMC Med. Genet.* 19, 196. <https://doi.org/10.1186/s12881-018-0709-3>.
58. Perli, E., Giordano, C., Tuppen, H.A.L., Montopoli, M., Montanari, A., Orlandi, M., Pisano, A., Catanzaro, D., Caparrotta, L., Musumeci, B., et al. (2012). Isoleucyl-tRNA synthetase levels modulate the penetrance of a homoplasmic m.4277T>C mitochondrial tRNA(Ile) mutation causing hypertrophic cardiomyopathy. *Hum. Mol. Genet.* 21, 85–100. <https://doi.org/10.1093/hmg/ddr440>.
59. Ferdinandusse, S., Friederich, M.W., Burlina, A., Ruiten, J.P.N., Coughlin, C.R., 2nd, Dishop, M.K., Gallagher, R.C., Bedoyan, J.K., Vaz, F.M., Waterham, H.R., et al. (2015). Clinical and biochemical characterization of four patients with mutations in ECHS1. *Orphanet J. Rare Dis.* 10, 79. <https://doi.org/10.1186/s13023-015-0290-1>.
60. Du, J., Li, Z., Li, Q.Z., Guan, T., Yang, Q., Xu, H., Pritchard, K.A., Camara, A.K.S., and Shi, Y. (2013). Enoyl coenzyme A hydratase domain-containing 2, a potential novel regulator of myocardial ischemia injury. *J. Am. Heart Assoc.* 2, e000233. <https://doi.org/10.1161/JAHA.113.000233>.
61. Sharpe, A.J., and McKenzie, M. (2018). Mitochondrial fatty acid oxidation disorders associated with short-chain enoyl-CoA hydratase (ECHS1) deficiency. *Cells* 7, 46. <https://doi.org/10.3390/cells7060046>.
62. Sharkia, I., Hadad Erlich, T., Landolina, N., Assayag, M., Motzik, A., Rachmin, I., Kay, G., Porat, Z., Tshori, S., Berkman, N., et al. (2017). Pyruvate dehydrogenase has a major role in mast cell function, and its activity is regulated by mitochondrial microphthalmia transcription factor. *J. Allergy Clin. Immunol.* 140, 204–214.e8. <https://doi.org/10.1016/j.jaci.2016.09.047>.
63. Guo, W., Ma, J., Yang, Y., Guo, S., Zhang, W., Zhao, T., Yi, X., Wang, H., Wang, S., Liu, Y., et al. (2020). ATP-citrate lyase epigenetically potentiates oxidative phosphorylation to promote melanoma growth and adaptive resistance to MAPK inhibition. *Clin. Cancer Res.* 26, 2725–2739. <https://doi.org/10.1158/1078-0432.CCR-19-1359>.
64. Haq, R., Shoag, J., Andreu-Perez, P., Yokoyama, S., Edelman, H., Rowe, G.C., Frederick, D.T., Hurley, A.D., Nellore, A., Kung, A.L., et al. (2013). Oncogenic BRAF regulates oxidative metabolism via PGC1alpha and MITF. *Cancer Cell* 23, 302–315. <https://doi.org/10.1016/j.ccr.2013.02.003>.
65. Ferguson, J., Smith, M., Zudaire, I., Wellbrock, C., and Arozarena, I. (2017). Glucose availability controls ATF4-mediated MITF suppression to drive melanoma cell growth. *Oncotarget* 8, 32946–32959. <https://doi.org/10.18632/oncotarget.16514>.
66. Holden, H.M., Rayment, I., and Thoden, J.B. (2003). Structure and function of enzymes of the Leloir pathway for galactose metabolism. *J. Biol. Chem.* 278, 43885–43888. <https://doi.org/10.1074/jbc.R300025200>.
67. Mueckler, M., and Thorens, B. (2013). The SLC2 (GLUT) family of membrane transporters. *Mol. Aspects Med.* 34, 121–138. <https://doi.org/10.1016/j.mam.2012.07.001>.
68. Hamberger, A., and Hyden, H. (1963). Inverse enzymatic changes in neurons and glia during increased function and hypoxia. *J. Cell Biol.* 16, 521–525. <https://doi.org/10.1083/jcb.16.3.521>.
69. Díaz-García, C.M., Meyer, D.J., Nathwani, N., Rahman, M., Martínez-François, J.R., and Yellen, G. (2021). The distinct roles of calcium in rapid control of neuronal glycolysis and the tricarboxylic acid cycle. *Elife* 10, e64821. <https://doi.org/10.7554/eLife.64821>.
70. Larson, P.E.Z., and Gordon, J.W. (2021). Hyperpolarized metabolic MRI-acquisition, reconstruction, and analysis methods. *Metabolites* 11, 386. <https://doi.org/10.3390/metabo11060386>.
71. Le Page, L.M., Guglielmetti, C., Taglang, C., and Chaumeil, M.M. (2020). Imaging brain metabolism using hyperpolarized (13)C magnetic resonance spectroscopy. *Trends Neurosci.* 43, 343–354. <https://doi.org/10.1016/j.tins.2020.03.006>.
72. Zimmer, E.R., Parent, M.J., Souza, D.G., Leuzy, A., Lecrux, C., Kim, H.I., Gauthier, S., Pellerin, L., Hamel, E., and Rosa-Neto, P. (2017). [(18)F] FDG PET signal is driven by astroglial glutamate transport. *Nat. Neurosci.* 20, 393–395. <https://doi.org/10.1038/nn.4492>.
73. Dayton, T.L., Gocheva, V., Miller, K.M., Israelsen, W.J., Bhutkar, A., Clish, C.B., Davidson, S.M., Luengo, A., Bronson, R.T., Jacks, T., and Vander Heiden, M.G. (2016). Germline loss of PKM2 promotes metabolic distress and hepatocellular carcinoma. *Genes Dev.* 30, 1020–1033. <https://doi.org/10.1101/gad.278549.116>.
74. Shin, B.C., Cepeda, C., Eghbali, M., Byun, S.Y., Levine, M.S., and Devasakar, S.U. (2021). Adult glut3 homozygous null mice survive to demonstrate neural excitability and altered neurobehavioral responses reminiscent of neurodevelopmental disorders. *Exp. Neurol.* 338, 113603. <https://doi.org/10.1016/j.expneurol.2021.113603>.
75. Goodpaster, B.H., and Sparks, L.M. (2017). Metabolic flexibility in health and disease. *Cell Metab.* 25, 1027–1036. <https://doi.org/10.1016/j.cmet.2017.04.015>.
76. Motori, E., Atanassov, I., Kochan, S.M.V., Folz-Donahue, K., Sakthivelu, V., Gialvalisco, P., Toni, N., Puyal, J., and Larsson, N.G. (2020). Neuronal metabolic rewiring promotes resilience to neurodegeneration caused by mitochondrial dysfunction. *Sci. Adv.* 6, eaba8271. <https://doi.org/10.1126/sciadv.aba8271>.
77. Tramunt, B., Smati, S., Grandgeorge, N., Lenfant, F., Arnal, J.F., Montagner, A., and Gourdy, P. (2020). Sex differences in metabolic regulation and diabetes susceptibility. *Diabetologia* 63, 453–461. <https://doi.org/10.1007/s00125-019-05040-3>.
78. Du, L., Hickey, R.W., Bayir, H., Watkins, S.C., Tyurin, V.A., Guo, F., Kochanek, P.M., Jenkins, L.W., Ren, J., Gibson, G., et al. (2009). Starving

- neurons show sex difference in autophagy. *J. Biol. Chem.* 284, 2383–2396. <https://doi.org/10.1074/jbc.M804396200>.
79. Christianto, A., Baba, T., Takahashi, F., Inui, K., Inoue, M., Suyama, M., Ono, Y., Ohkawa, Y., and Morohashi, K.I. (2021). Sex differences in metabolic pathways are regulated by Pfkfb3 and Pdk4 expression in rodent muscle. *Commun. Biol.* 4, 1264. <https://doi.org/10.1038/s42003-021-02790-y>.
 80. Levéen, P., Kotarsky, H., Mörgelin, M., Karikoski, R., Elmér, E., and Fellman, V. (2011). The GRACILE mutation introduced into Bcs1l causes post-natal complex III deficiency: a viable mouse model for mitochondrial hepatopathy. *Hepatology* 53, 437–447. <https://doi.org/10.1002/hep.24031>.
 81. Kotarsky, H., Keller, M., Davoudi, M., Levéen, P., Karikoski, R., Enot, D.P., and Fellman, V. (2012). Metabolite profiles reveal energy failure and impaired beta-oxidation in liver of mice with complex III deficiency due to a BCS1L mutation. *PLoS One* 7, e41156. <https://doi.org/10.1371/journal.pone.0041156>.
 82. Purhonen, J., Grigorjev, V., Ekiert, R., Aho, N., Rajendran, J., Pietras, R., Truvé, K., Wikström, M., Sharma, V., Osyczka, A., et al. (2020). A spontaneous mitonuclear epistasis converging on Rieske Fe-S protein exacerbates complex III deficiency in mice. *Nat. Commun.* 11, 322. <https://doi.org/10.1038/s41467-019-14201-2>.
 83. Colville, C.A., Seatter, M.J., Jess, T.J., Gould, G.W., and Thomas, H.M. (1993). Kinetic analysis of the liver-type (GLUT2) and brain-type (GLUT3) glucose transporters in *Xenopus* oocytes: substrate specificities and effects of transport inhibitors. *Biochem. J.* 290, 701–706. <https://doi.org/10.1042/bj2900701>.
 84. Maher, F., Davies-Hill, T.M., and Simpson, I.A. (1996). Substrate specificity and kinetic parameters of GLUT3 in rat cerebellar granule neurons. *Biochem. J.* 315, 827–831. <https://doi.org/10.1042/bj3150827>.
 85. Reagan, L.P., Rosell, D.R., Alves, S.E., Hoskin, E.K., McCall, A.L., Charon, M.J., and McEwen, B.S. (2002). GLUT8 glucose transporter is localized to excitatory and inhibitory neurons in the rat hippocampus. *Brain Res.* 932, 129–134. [https://doi.org/10.1016/S0006-8993\(02\)02308-9](https://doi.org/10.1016/S0006-8993(02)02308-9).
 86. Simpson, I.A., Chundu, K.R., Davies-Hill, T., Honer, W.G., and Davies, P. (1994). Decreased concentrations of GLUT1 and GLUT3 glucose transporters in the brains of patients with Alzheimer's disease. *Ann. Neurol.* 35, 546–551. <https://doi.org/10.1002/ana.410350507>.
 87. Manzo, E., Lorenzini, I., Barrameda, D., O'Conner, A.G., Barrows, J.M., Starr, A., Kovalik, T., Rabichow, B.E., Lehmkuhl, E.M., Shreiner, D.D., et al. (2019). Glycolysis upregulation is neuroprotective as a compensatory mechanism in ALS. *Elife* 8, e45114. <https://doi.org/10.7554/eLife.45114>.
 88. Horlbeck, M.A., Gilbert, L.A., Villalta, J.E., Adamson, B., Pak, R.A., Chen, Y., Fields, A.P., Park, C.Y., Corn, J.E., Kampmann, M., and Weissman, J.S. (2016). Compact and highly active next-generation libraries for CRISPR-mediated gene repression and activation. *Elife* 5, e19760. <https://doi.org/10.7554/eLife.19760>.
 89. Chen, E.Y., Tan, C.M., Kou, Y., Duan, Q., Wang, Z., Meirelles, G.V., Clark, N.R., and Ma'ayan, A. (2013). Enrichr: interactive and collaborative HTML5 gene list enrichment analysis tool. *BMC Bioinformatics* 14, 128. <https://doi.org/10.1186/1471-2105-14-128>.
 90. Kuleshov, M.V., Jones, M.R., Rouillard, A.D., Fernandez, N.F., Duan, Q., Wang, Z., Koplev, S., Jenkins, S.L., Jagodnik, K.M., Lachmann, A., et al. (2016). Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Res.* 44, W90–W97. <https://doi.org/10.1093/nar/gkw377>.
 91. Xie, Z., Bailey, A., Kuleshov, M.V., Clarke, D.J.B., Evangelista, J.E., Jenkins, S.L., Lachmann, A., Wojciechowicz, M.L., Kropiwnicki, E., Jagodnik, K.M., et al. (2021). Gene set knowledge discovery with Enrichr. *Curr. Protoc.* 1, e90. <https://doi.org/10.1002/cpz1.90>.
 92. Berthet, A., Margolis, E.B., Zhang, J., Hsieh, I., Zhang, J., Hnasko, T.S., Ahmad, J., Edwards, R.H., Sesaki, H., Huang, E.J., and Nakamura, K. (2014). Loss of mitochondrial fission depletes axonal mitochondria in midbrain dopamine neurons. *J. Neurosci.* 34, 14304–14317. <https://doi.org/10.1523/JNEUROSCI.0930-14.2014>.
 93. Johnson, E.C.B., Ho, K., Yu, G.Q., Das, M., Sanchez, P.E., Djukic, B., Lopez, I., Yu, X., Gill, M., Zhang, W., et al. (2020). Behavioral and neural network abnormalities in human APP transgenic mice resemble those of App knock-in mice and are modulated by familial Alzheimer's disease mutations but not by inhibition of BACE1. *Mol. Neurodegener.* 15, 53. <https://doi.org/10.1186/s13024-020-00393-5>.
 94. Arguello, A., Meisner, R., Thomsen, E.R., Nguyen, H.N., Ravi, R., Simms, J., Lo, I., Speckart, J., Holtzman, J., Gill, T.M., et al. (2021). Iduronate-2-sulfatase transport vehicle rescues behavioral and skeletal phenotypes in a mouse model of Hunter syndrome. *JCI Insight* 6, e145445. <https://doi.org/10.1172/jci.insight.145445>.
 95. Xia, Z., and Liu, Y. (2001). Reliable and global measurement of fluorescence resonance energy transfer using fluorescence microscopes. *Biophys. J.* 81, 2395–2402. [https://doi.org/10.1016/S0006-3495\(01\)75886-9](https://doi.org/10.1016/S0006-3495(01)75886-9).
 96. Shields, L.Y., Li, H., Nguyen, K., Kim, H., Doric, Z., Garcia, J.H., Gill, T.M., Haddad, D., Vossel, K., Calvert, M., and Nakamura, K. (2021). Mitochondrial fission is a critical modulator of mutant APP-induced neural toxicity. *J. Biol. Chem.* 296, 100469. <https://doi.org/10.1016/j.jbc.2021.100469>.

STAR★METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Rabbit monoclonal anti-PKM1	Cell signaling	Cat #7067; RRID: AB_2715534
Rabbit monoclonal anti-PKM2	Novus	Cat# NBP1-48308; RRID: AB_10011057
Chicken monoclonal anti-MAP2	Abcam	Cat# ab5392; RRID: AB_2138153
Mouse monoclonal anti-NeuN	Millipore	Cat# MAB377; RRID: AB_2298772
Rabbit monoclonal anti-NeuN	Abcam	Cat# ab177487; RRID: AB_2532109
Rabbit anti-BCS1L	Novus	Cat# NBP2-92916
Rabbit anti-Mitf	Abcam	Cat# ab122982; RRID: AB_10902226
Anti-rabbit IgG Alexa Fluor 488	Thermo Fisher Scientific	A11034; RRID: AB_2576217
Anti-mouse IgG Alexa Fluor 594	Thermo Fisher Scientific	A-11037; RRID: AB_2534095
Anti-chicken IgG Alexa Fluor 350	Thermo Fisher Scientific	PI62272
Hoechst	Invitrogen	H1399
Bacterial and virus strains		
Lentivirus (2 ND gen) PKM sgRNA	UCSF Viracore	This study
Lentivirus (2 ND gen) GLUT3 sgRNA	UCSF Viracore	This study
Lentivirus (2 ND gen) NTG sgRNA	UCSF Viracore	This study
Chemicals, peptides, and recombinant proteins		
Paraformaldehyde	Electron Microscopy Science	Cat# 30525-89-4
HEPES	Sigma Aldrich	Cat #3375
D-Glucose	Sigma Aldrich	Cat# G7021
Sodium Pyruvate	Sigma Aldrich	Cat# P2262
Cytochalasin B	Sigma Aldrich	Cat# C2743
2-Deoxy-D-glucose	Sigma Aldrich	Cat# D8375
Rotenone	Sigma Aldrich	r8875-1g
Iodoacetic Acid	Thermo Fisher Scientific	Cat# 35603
Sucrose	Sigma Aldrich	Cat# 9378
Phosphate buffered saline	UCSF Media Production	CCFAL001
Isopentane	Sigma Aldrich	Cat# PHR1661
Triton X-100	Sigma Aldrich	Cat# T8787
Vectashield antifade mounting medium	Vector Laboratories	Cat# H-1000-10
Prolong gold antifade mountant	Thermo Fisher Scientific	Cat# P36941
Prolong diamond antifade mountant	Thermo Fisher Scientific	Cat# P36970
Goat serum	Thermo Fisher Scientific	Cat# 16210064
MEM Eagle's with Earle's BSS Medium	UCSF Cell Culture Facility	CCFAC001
Fetal bovine serum	Hyclone	SH3007103
Glutamax	Gibco	Cat# 175007
Serum extender	Thermo Fisher Scientific	355006
Uridine	Sigma Aldrich	Cat# U3750
5-fluoro-2-deoxycytidine	Sigma Aldrich	Cat# F0503
[U- ¹³ C]glucose	Cambridge Isotope Labs	CLM-1396-1
[U- ¹³ C]Galactose	Sigma Aldrich	Cat# 605379
Neurobasal A media	Gibco	A2477501

(Continued on next page)

<i>Continued</i>		
REAGENT or RESOURCE	SOURCE	IDENTIFIER
NEAA	Gibco	11140050
N2 supplement	Gibco	17502048
B27 supplement	Gibco	17504044
GlutaMAX	Gibco	35050061
NT-3	Stemcell Tech	78074
BDNF	Stemcell Tech	78005
Laminin	Gibco	23017015
Ammonium acetate	Sigma Aldrich	Cat# 32301
Methanol	Sigma Aldrich	Cat# 34860
StemFlex medium	ThermoFisher	A3349401
Matrigel	Corning	Cat# 356234
DMEM/F12	Gibco	Cat#12660012
NEAA	Gibco	Cat#11140050
Doxycycline hyclate	Sigma Aldrich	D9891
ROCK Inhibitor Y27632	Stemcell Tech	72302
[U- ¹³ C] glucose	Cambridge Isotope Labs	CLM-1396-1
Gadavist	Bayer	768511
[1- ¹³ C]pyruvate	Sigma-Aldrich	677175
[¹⁸ F]-fluorodeoxyglucose	University of California San Francisco Radiopharmaceutical Facility	N/A
<i>Critical commercial assays</i>		
Glucometer	Ermaline Laboratories	Cat# 37321
Visium spatial gene expression slide & reagent kit	10X Genomics	1000184
Dual index Kit TT Set A	10X Genomics	1000215
Lonza electroporation Kit	Amaxa	Cat# VVSI-1001
Optimal Cutting Temperature	Tissue-Tek	Cat# 62550-12
<i>Experimental models: Cell lines</i>		
Mouse primary neuronal culture	N/A	This study
Human: WTC-11 + NGN2 + dCAS9-KRAB	Martin Kampmann UCSF.	Tian et al. ²²
<i>Experimental models: Organisms/strains</i>		
GLUT3 ^{lox/lox}	Jeff Rathmell, Vanderbilt	Contat et al. ²⁴
PKM1 ^{lox/lox}	Mathew Vander Heiden, MIT	Davidson et al. ⁴⁷
CamKII α	Jackson Laboratories	005359
<i>Oligonucleotides</i>		
PKM sgRNA GCCGCAGCGGGATAACCTT	IDT	Horlbeck et al. ⁸⁸
GLUT3 sgRNA GGTAAGACTTTGGATCCTTCCTG	IDT	Horlbeck et al. ⁸⁸
NTG sgRNA GCTGCATGGGGCGGAATCA	IDT	Semple et al. ³⁵ Horlbeck et al. ⁸⁸
<i>Recombinant DNA</i>		
iGlucoSnFR-mRuby	Jonathan Marvin, HHMI	N/A
ATP fluorescence resonance energy sensor	N/A	Mendelsohn et al. ⁵⁶
<i>Software and algorithms</i>		
Prism 8.4.3	GraphPad Software	https://www.graphpad.com/scientific-software/prism/
MetaMorph 7.7.3.0	Universal Imaging	RRID: SciRes_000136
SIVIC software	N/A	http://sourceforge.net/apps/trac/sivic/
MATLAB R2011b	The MathWorks Inc	N/A

(Continued on next page)

Continued

REAGENT or RESOURCE	SOURCE	IDENTIFIER
AMIRA software	Mercury Computer systems	N/A
ImageJ	NIH	http://imagej.nih.gov/ij
VivoQuant 4.0 software	Invicro	N/A
Dynamic Susceptibility Contrast MR Analysis	DSCoMAN	version 1.0
ANTS	N/A	https://github.com/ANTsX/ANTs
pClamp10 software	Molecular Devices	N/A
IgorPro6 software	Wavemetrics Inc	N/A
10X Space Ranger software	10X Genomics	https://support.10xgenomics.com/spatial-gene-expression/software/downloads/latest
10X Loupe Browser software	10X Genomics	https://www.10xgenomics.com/products/loupe-browser/downloads
Enrichr webtool	Avi Ma'ayan Lab	Chen et al. ⁸⁹ Kuleshov et al. ⁹⁰ Xie et al. ⁹¹

Other

LSM 880 Confocal Laser Scanning Microscope with Airyscan	Zeiss	N/A
SM2000 R microtome	Leica	N/A
Labconco CentriVap Concentrator	Labconco	N/A
Video tracking	Tracker	N/A
14.1 T vertical MR system	Agilent Technologies	N/A
Hypersense DNP polarizer	Oxford Instruments	N/A
PET/computerize tomography (CT) scanner	Siemens Healthcare	Inveon
COBRA Feldkamp reconstruction algorithm	Exxim Computing Corporation	N/A
VT1200 Vibratome	Leica	N/A
BSK5 vapor interface holding chamber	Scientific Systems Design Inc. Canada	N/A
RC-27LD submerged chamber	Warner Instruments	N/A
fDS2A-MKII voltage isolated stimulator	Digitimer North America	N/A
Ti-E inverted microscope	Nikon	N/A
iXon EMCCD camera	Andor Technology	N/A
VC-8 Perfusion valve system	Warner Instruments	N/A
A385 Current isolator	World Precision Instruments	N/A
SYS-A310 Accupulser signal generator	World Precision Instruments	N/A
Aperio Versa slide scanner	Leica	N/A

RESOURCE AVAILABILITY

Lead contact

Further information and requests for resources and reagents should be directed to Ken Nakamura (ken.nakamura@gladstone.ucsf.edu).

Materials availability

Information for resources and reagents are available from the [lead contact](#) upon request.

Data and code availability

- Visium datasets are available in [Tables S3](#) and [S4](#). Metabolomics data sets are available in [Tables S5](#), [S6](#) and [S7](#).
- This paper does not report original code.
- Further information and requests for reagents should be directed to and will be fulfilled by the [lead contact](#), Ken Nakamura (ken.nakamura@ucsf.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Mice

All animal experimental procedures were conducted in accordance with the Guide for the Care and Use of Laboratory Animals, as adopted by the National Institutes of Health, and with approval of the University of California, San Francisco Institutional Animal Care and Use Committee. Animals used in this project were up to 20 months of age. All mice were housed in a state-of-the-art barrier facility managed by the UCSF Laboratory Animal Resource Center (LARC). Animal care and use in this research are covered under the UCSF “Assurance of Compliance with PHS Policy on Humane Care and Use of Laboratory Animals by Awardee Institutions” number A3400-01. Experiments were performed on age-matched mice of either sex, with the specific number of males and females matched between groups in each experiment (typically divided (50:50 between males and females). All mouse lines were maintained on a C57Bl/6 background (The Jackson Laboratory; RRID:IMSR_JAX:000664).

Generation of iPSC lines and iPSC-derived neuronal cultures

Human WTC-11 induced pluripotent stem cells expressing both doxycycline inducible neurogenin 2 (Ngn2) and inducible CRISPRi (dCas9-KRAB) machinery were received from Martin Kampmann.²² Lentivirus expressing sgRNA targeting PKM sgRNA sequence GCCGCAGCGGGATAACCTT) or GLUT3 (sgRNA sequence GGTAAGACTTTGGATCCTTCCTG), or non-targeting control (GCTGCATGGGGCGCAATCA) were produced by the UCSF Viracore. iPSCs were incubated with lentivirus for 24h with 8 μ g/mL polybrene. After two days of culture in StemFlex medium (ThermoFisher A3349401), transfected iPSCs were selected with 1 μ g/mL puromycin for 1 day.

iPSCs were cultured in StemFlex media in a 0.1 mg/mL Matrigel (Corning 356234) coated culture flask. Differentiation of the iPSCs was divided into 2 parts: pre-differentiation and maturation. Pre-differentiation consisted of 3 days of culture in pre-differentiation media and maturation consisted of 11 days in neuronal differentiation media. Pre-differentiation media contained Knockout DMEM/F12 (Gibco 12660012), 1X NEAA (Gibco 11140050), 1X N2 supplement (Gibco 17502048), 10 ng/mL NT-3 (Stemcell Tech 78074), 10 ng/mL BDNF (Stemcell Tech 78005), and 1 μ g/mL of laminin (Gibco 23017015). On the first day of differentiation, iPSCs were passaged onto a 0.1 mg/mL Matrigel coated culture flask with pre-differentiation media containing doxycycline (2 μ g/mL) and ROCK Inhibitor (10 nM). The pre-differentiation media was then refreshed on the third day of pre-differentiation. On the fourth day of differentiation, pre-differentiated neurons were passaged into 5 μ g/mL laminin coated 6-well plates at a density of 500,000 cells/well with the neuronal differentiation media; 20 μ M of trimethoprim was used to induce the target knockdowns (PKM or GLUT3) 2 days after the transition to the neuronal differentiation media.²² Neuronal differentiation media contained 1:1 of DMEM/F12 (Gibco 11320033) and Neurobasal A media (Gibco 10888022), 1X NEAA (Gibco 11140050), 0.5X N2 supplement (Gibco 17502048), 0.5X B27 supplement (Gibco 17504044), 0.5X GlutaMAX (Gibco 35050061), 10 ng/mL NT-3 (Stemcell Tech 78074), 10 ng/mL BDNF (Stemcell Tech 78005), and 1 μ g/mL of laminin (Gibco 23017015). The neuronal differentiation media was refreshed once a week to feed the neurons.

Primary neuronal cultures

Primary hippocampal cultures were dissected from early postnatal (P0 to P1) mouse pups.^{30,92} Neurons were transfected using electroporation (Lonza, VVSI-1001 with the relevant plasmids and cultured in MEM Eagle's with Earle's BSS Medium (UCSF Cell Culture Facility, CCFAC001) supplemented with 5% fetal bovine serum (Hyclone, SH3007103), 1% Glutamax (Gibco 175007), 21mM glucose, 2% B27 supplement (Gibco 17504-044), and 0.1% serum extender (Fisher 355006). Neurons were plated at 600–900 cells/mm², and uridine (10 μ M, Sigma U3003) and 5-fluoro-2-deoxycytidine (10 μ M, Sigma F0503) were added on day 4–5 to minimize glial overgrowth. Neurons were cultured at 37°C with 5% CO₂ for 9–11 days before live imaging or fixation.

METHOD DETAILS

Metabolomics

Human iPSC-derived neurons were matured for a total of 12 days, including a 24 h incubation with media containing a [U-¹³C] metabolic probe prior to metabolite collection. The base media for targeted metabolomics contained Neurobasal A media without glucose or pyruvate (Gibco A2477501), 1X NEAA (Gibco 11140050), 0.5X N2 supplement (Gibco 17502048), 0.5X B27 supplement (Gibco 17504044), 0.5X GlutaMAX (Gibco 35050061), 10 ng/mL NT-3 (Stemcell Tech 78074), 10 ng/mL BDNF (Stemcell Tech 78005), and 1 μ g/mL of laminin (Gibco 23017015). The media was supplemented with either [U-¹³C]glucose (Cambridge Isotope Labs CLM-1396-1) or [U-¹³C]galactose (Sigma-Aldrich 605379) for targeted metabolomics. To extract cellular metabolites, cells were washed in ammonium acetate (150 mM, pH 7.3) at 4C, and then incubated with 80% methanol at –80C for 20 min, prior to centrifugation at 14,000 g for 15 min. To extract cell media metabolites, the ¹³C-labeled media contained in each well post-incubation was collected and spun at 14,000 g for 10 min. The clarified media was then added to 80% methanol chilled to 4C, followed up with a 1 h incubation at –80C. Media samples were then centrifuged at 14,000 g for 10 min post incubation. Metabolite supernatants were dried in a Labconco CentriVap prior to storage at –80C. Sample metabolites were quantified at the UCLA Metabolomics Core. Metabolic media conditions are detailed below.

Condition	Metabolic probe	Description
1	[U- ¹³ C]glucose 20.5 mM	Differentiation media with basal level of glucose
2	[U- ¹³ C]glucose 1.5 mM	Physiologically relevant brain glucose level
3	[U- ¹³ C]glucose 1.5 mM	1:1 glucose:galactose media tracing glucose
4	[U- ¹³ C]galactose 1.5 mM	Galactose replacing all glucose from condition 2

Galactose media measurements

A colorimetric galactose assay kit (Sigma Aldrich MAK012) was used to measure the galactose content of neuronal differentiation media supplement B27 (Gibco 17504044) using the protocol included with the kit. Samples were measured using a Molecular Devices Microplate Reader at an absorbance of 570 nm.

Behavioral testing

Elevated plus maze

The elevated plus maze (Kinder Scientific) consists of two open arms (without walls) and two closed arms (with walls) elevated 63 cm above the floor. Mice were acclimated to a dimly-lit testing room for 1 h prior to testing and mice were allowed to freely explore the maze for 10 min. The time, distance, and entries into the open and closed arms are calculated based on infrared photobeam breaks.

Open field

Spontaneous locomotor activity was assessed in the open field test (Flex-Field, San Diego Instruments) using standard procedures.⁹³ Mice were placed in the center of a clear acrylic chamber and allowed to freely explore for 15 min. The photobeams that line the perimeter of the chamber detect horizontal and vertical movements. Ambulatory and fine movements are separated recorded. Fine movements are defined as breaking 2 beams consecutively and ambulatory movements are defined as breaking 3 beams or more.

Active place avoidance

Visual-spatial learning and memory was measured by the active place avoidance task (BioSignal Corp).⁹⁴ Mice were trained to a new aversive zone for each time point. A habituation trial was conducted on the first day to determine baseline activity, followed by 3 days of shock training to the aversive zone. A probe and reinstatement trial was conducted 24 h after the last day of training. The number of entries, shocks, and the maximum time of avoidance to the aversive zone are calculated based on the video tracking software (Tracker).

Hot plate

Nociception was measured on the hot plate test (IITC). The surface of the aluminum hot plate was heated to 52°C and mice were placed in an enclosure atop the hot plate. The response latency is determined by a hindpaw lick, flick, or jump. The maximum latency was 30 s to prevent injury.

Histology and immunostaining

For *in vivo* histology, mice were perfused with phosphate-buffered saline (PBS) and then 4% paraformaldehyde (PFA). Brains were postfixed in 4% PFA overnight, and cryoprotected in 30% sucrose. Brain sections were set at 30–40 μm and prepared using a sectioning microtome (Leica SM2000 R). For immunofluorescence, brain sections were washed 3X in PBS, blocked 1 h at room temperature with blocking buffer (5% goat serum (Abcam, Cat #ab7481), 0.2% Triton in PBS), and incubated with primary antibodies at 4°C overnight with blocking buffer. The following antibodies were used: rabbit anti-PKM1 (1:400; Cell signaling Cat #7067, RRID: AB_2715534); rabbit anti-PKM2 (1:400; Novus Cat# NBP1-48308, RRID AB_10011057); chicken anti-MAP2 (1:1000; Abcam Cat# ab5392, RRID:AB_2138153); mouse anti-NeuN (1:1000; Millipore Cat# MAB377, RRID:AB_2298772); rabbit anti-NeuN (1:1000; Abcam Cat# ab177487, RRID: AB_2532109); rabbit anti-BCS1L (1:1000; Novus Cat# NBP2-92916); rabbit anti-Mitf (1:1000; Abcam Cat# ab122982, RRID: AB_10902226). Sections were rinsed 3X in PBS and incubated 2h at RT with the following secondary antibodies: Alexa Fluor 350, 488, 594, or 647 anti-mouse, chicken, or rabbit IgG (1:500; Invitrogen). Sections were imaged on a Zeiss LSM880 confocal microscope with Airyscan detector.

Neuronal density was calculated by dividing the total count number of NeuN and DAPI positive cells within CA1 region by the CA1 region area. Quantification of the density was performed blind to genotype with MetaMorph software (version 7.7.3.0; Universal Imaging, RRID:SciRes_000136).

In vivo MR acquisitions

All *in vivo* MR experiments were conducted on a 14.1 T vertical MR system (Agilent Technologies, Palo Alto, CA) equipped with 100 G/cm gradients and a dual tune ¹H-¹³C volume coil (Ø₁ = 40 mm, for HP ¹³C MRSI and T₂-weighted MRI) or a single tuned ¹H proton coil (Ø₁ = 40 mm, for dynamic susceptibility contrast (DSC) MRI and T₂-weighted MRI). For each imaging session, mice were anesthetized using isoflurane (2% in O₂) and a 27 G catheter was secured in the tail vein to allow for intravenous (iv) injection of the HP

probe or gadolinium contrast agent. Animals were then positioned in a dedicated cradle under constant anesthesia and placed in the MR bore; respiration was continuously monitored during all acquisitions to ensure animal well-being. First, axial T_2 -weighted images were acquired for adequate positioning of the grid used for HP ^{13}C acquisitions. The following parameters were used: echo time (TE)/repetition time (TR) = 20/1200 m, slice thickness = 1.8 mm, number of averages = 2, matrix = 256 × 256, field of view (FOV) = 30 × 30 mm². For HP ^{13}C MRSI acquisitions, 24 μL of [1- ^{13}C]pyruvate preparation, was hyperpolarized using a Hypersense DNP polarizer (Oxford Instruments) for one approximately hour.⁴⁰ After polarization, HP [1- ^{13}C]pyruvate was rapidly dissolved in isotonic buffer (pH~7) to a final concentration of 80 mM. A final volume of 300 μL of the HP [1- ^{13}C]pyruvate solution was then injected iv over 12 sec through the tail vein catheter. 2D ^{13}C MRSI data were acquired 18 s after injection of the HP [1- ^{13}C]pyruvate solution, using the following parameters: TE/TR = 0.56/68 m; spectral width = 4006 Hz; 256 points; flip angle (FA) = 10°; FOV = 30 × 30 mm²; 5 mm slice thickness. Next, axial T_2 -weighted Fast Spin Echo images were acquired for the evaluation of brain volume, using the following parameters: TE/TR = 12/2000 m, echo train = 8, slice thickness = 0.5 mm, number of averages = 8, matrix = 256 × 256, FOV = 30 × 30 mm². For DSC MR images, a high contrast single slice axial T_2 -weighted Fast Spin Echo image was acquired using the following parameters: TE/TR = 30/3000 m, slice thickness = 2 mm, matrix = 128 × 128, FOV = 20 × 20 mm². Next, a series of single slice axial T_2 -weighted Fast Spin Echo images were acquired, using the following parameters: TE/TR = 30/1000 m, slice thickness = 2 mm, matrix = 128 × 128, FOV = 20 × 20 mm², 150 repetitions. In order to obtain a pre-contrast baseline, the first 25 s of the dynamic MR acquisition were performed before the administration of the contrast agent (Gadavist, Bayer). At 25 s, an iv bolus of 1 mmol/kg of body weight of Gadavist was delivered through the tail vein catheter.

Ex vivo MRI acquisitions

Mice were euthanized and perfused with a phosphate buffered saline (PBS) solution followed by a 4% paraformaldehyde (PFA) solution. *Ex vivo* MR experiments were conducted on a 14.1 T vertical MR system (Agilent Technologies, Palo-Alto, CA) equipped with a single tuned ^1H proton coil ($\Phi_1 = 20$ mm). *Ex vivo* specimens were submerged in a Fluorinert (Sigma-Aldrich) solution and high-resolution 3D gradient echo images were acquired from the intact brain left in the skull using the following parameters: TE/TR = 15/75 m, number of averages = 6, matrix = 256 × 256 × 256, FOV = 12.8 × 12.8 × 12.8mm².

MR data analysis

HP ^{13}C MRSI datasets were analyzed using the SIVIC software (<http://sourceforge.net/apps/trac/sivic/>) and custom-built programs written in MATLAB (MATLAB R2011b, The MathWorks Inc.). The k-space dimensions were zero-filled by a factor of two resulting in a 32 × 32 matrix. The area under the curve (AUC) of HP [1- ^{13}C]pyruvate and AUC of HP [1- ^{13}C]lactate lorentzian fits were measured and the corresponding HP [1- ^{13}C]lactate-to-pyruvate ratio was calculated for each voxel. Regions of interest (ROIs) including the hippocampus and the thalamic area were defined on T_2 -weighted images and HP [1- ^{13}C]lactate-to-pyruvate ratios were reported as mean of the corresponding voxels. For analyses of *in vivo* MRI data, ROIs were manually delineated on T_2 -weighted images according to the Franklin and Paxinos anatomical mouse brain atlas using AMIRA software (Mercury Computer systems, San Diego, USA) and the corresponding ROI volume calculated. Specifically, the ROIs consisted of the entire brain, the ventricles and the hippocampus. In a similar manner, for analyses of *ex vivo* anatomical MRI, ROIs were manually delineated over the entire brain, the hippocampus and the CA1 regions, and the corresponding volume calculated for each ROI. For analyses of DSC images, Dynamic Susceptibility Contrast MR Analysis (DSCoMAN, version 1.0) plugin of ImageJ 1.52p (<http://imagej.nih.gov/ij>) was used to create cerebral blood volume (CBV) and cerebral blood flow (CBF) maps. ROIs including the hippocampus, thalamic areas and corpus callosum were manually delineated on the high contrast T_2 -weighted image acquired prior to the DSC acquisitions. These ROIs were then propagated onto the CBV and CBF maps and mean relative CBV and CBF values were obtained. In order to compare animals and groups, CBV and CBF values of the hippocampus and thalamic areas were normalized to the values obtained in the corpus callosum.

In vivo PET acquisitions

All scans were performed on a dedicated small animal PET/computerized tomography (CT) scanner (Inveon, Siemens Healthcare, Malvern, PA, USA). Mice received 71 ± 4.5 μCi of [^{18}F]-fluorodeoxyglucose (FDG) in 0.1 mL volume iv via tail vein. Fifty-five minutes after radiotracer administration, static PET images were collected (10-min acquisition time), followed by a 10-min CT acquisition for attenuation correction of PET reconstruction. Animals were kept awake at room temperature during the 55 min uptake time. The animals were placed under anesthesia (2% isoflurane in O_2) during the injections and scans.

PET data analysis

PET images were reconstructed using the ordered subsets expectation maximization (OSEM) algorithm, provided by the manufacturer. The resulting PET images had a 128 × 128 × 159 matrix with a voxel size of 0.776 × 0.776 × 0.796 mm³. CT images were reconstructed using a cone-beam Feldkamp reconstruction algorithm (COBRA, Exxim Computing Corporation, Pleasanton, CA, USA). The reconstructed CT images had a 512 × 512 × 662 matrix size with an isotropic voxel size of 0.191 mm³. The co-registered attenuation map from CT, obtained via a pre-derived rigid transformation matrix, was used for attenuation correction of the PET data. MR anatomical templates were created using the open source registration toolkit ANTS (<https://github.com/ANTsX/ANTs>) for each genotype and sex using *in vivo* T_2 -weighted images, leading to the creation of 8 MR templates: female GLUT3 WT, female GLUT3 KO, male GLUT3 WT, male GLUT3 KO, female PKM1 WT, female PKM1 KO, male PKM1 WT, male PKM1 KO. The PET/CT images of each

individual mouse was co-registered to the corresponding MR template using VivoQuant 4.0 software (Invivo). Hippocampus and CA1 ROIs were manually delineated using the MR template and corresponding mean percent-injected dose per grams (%ID/g) values were obtained.

Electrophysiology

Animals were anesthetized with isoflurane and decapitated. The brains were rapidly removed from the skull and placed in the ice-cold (2–5°C) slicing solution. Slicing solution contained (in mM): 110 choline chloride, 2.5 KCl, 26 NaHCO₃, 10 MgCl₂, 1.25 NaH₂PO₄, 0.5 CaCl₂, 10 glucose, 3 Na Pyruvate, 1 L-Ascorbic acid, pH 7.4.

300 μm-thick sagittal slices were cut from both hemispheres using a Vibratome (VT1200, Leica) and transferred to a 95% O₂-CO₂ vapor interface holding chamber (BSK5, Scientific Systems Design Inc. Canada) containing artificial cerebrospinal fluid (ACSF) where they were allowed to recover at 34°C for 1 h and held at room temperature (20–22°C) afterward. ACSF contained (in mM): 126 NaCl, 2.5 KCl, 1.5 CaCl₂, 1.5 MgCl₂, 26 NaHCO₃, 1.25 NaH₂PO₄, 10 glucose, and 1.5 L-Ascorbic acid, pH 7.4. For recordings, slices were transferred to a submerged chamber (RC-27LD, Warner Instruments, USA) where they were continuously superfused on both sides with oxygenated ACSF at 32 °C at a flow rate of 10 mL/min.

Local Field Potentials (LFPs) were elicited by orthodromic stimulation of Schaffer collaterals by concentric bipolar stimulating electrode (FHC, Inc., USA) connected to a constant voltage isolated stimulator (DS2A-MKII, Digitimer North America) and placed in CA2 str. radiatum. Repetitive network activity was induced by a 300-pulse 10Hz stimulus train delivered to Schaffer collaterals; stimulus intensity was chosen to elicit a str. Pyramidale population spike amplitude at 50% of maximum. LFPs were recorded with a glass borosilicate microelectrodes filled with ACSF and placed in CA1 str. Radiatum (dendritic post-synaptic potentials, fPSPs), with the resulting LFP population response simultaneously in str. Pyramidale. Signals were sampled and digitized by MultiClamp 700B amplifier and Digidata 1550 acquisition system with pClamp10 software (Molecular Devices, USA), and analyzed using IgorPro6 software (Wavemetrics Inc., USA) running custom macros. fPSP slopes were analyzed as the linear fit slope values between 10% and 90% of fPSP peak. Input-output relationships were recorded as the fPSP slope values in response to increasing stimulation intensity (5–25mV), as well as the LFP population spike integral vs. fPSP slopes.

Live imaging of glucose or ATP

Live imaging was performed in Tyrode's buffer (127mM NaCl, 10mM HEPES, 30mM glucose, 10 mM pyruvate, 2.5 mM KCl, 2mM CaCl₂ and 2mM MgCl₂, pH 7.4) at room temperature with a 40X objective on a Nikon Ti-E inverted microscope with an iXon EMCCD camera (Andor Technology, Belfast, UK) and a perfusion valve system (VC-8, Warner Instruments, Hamden, CT, USA) controlled by MetaMorph Software. Field stimulations (5 Hz, 300s) were performed with an A385 current isolator and an SYS-A310 accupulser signal generator (World Precision Instruments).

For imaging experiments using the cytosolic glucose sensor (iGlucoSnFR-mRuby expressed under the synapsin-1 promoter, kind gift of Jonathan Marvin, HHMI) sequential images were taken from FITC (490/20 ex, 535/50 em) and Texas Red (572/35 ex, 632/60 em) channels (Chroma, 89014). The glucose sensor fluorescence was normalized to mRuby fluorescence to control for expression level. Total glucose uptake was blocked with cytochalasin B (10 μM, Sigma-Aldrich). For measurements with the ATP fluorescence resonance energy sensor (Mendelsohn et al., 2018), sequential images were taken from mClover (490/20 ex, 535/50 em), mApple (572/35 ex, 632/60 em), and FRET channels (490/20 ex, 632/60 em). Synaptic boutons were identified based on co-localization with BFP-synaptophysin. The FRET/donor ratio was calculated for each bouton following the formula: $FRET = (I_{FRET} - I_{Clover} * BT_{Clover} - I_{mApple} * BT_{mApple}) / I_{Clover}$. I_x is the background-corrected fluorescence intensity measured in that given channel. BT_{Clover} (donor bleed through) and BT_{mApple} (direct excitation of the acceptor) were calculated by expressing clover and mApple individually and determining the ratios of I_{FRET}/I_{Clover} and I_{FRET}/I_{mApple} independently.^{95,96}

Blood glucose level measurement

Blood samples were collected under non-fasted condition and obtained from the tail tip. The blood glucose level was measured using a glucometer (Cat #37321, Ermaine Laboratories).

Spatial gene expression

Spatial transcriptomics were acquired with Visium spatial gene expression kits (10X Genomics 1000184, 1000193, 1000215). Sample preparation, sample imaging, and library generation were completed in accordance with 10X Spatial Gene Expression protocols included with the kit. Briefly, fresh brain tissue was flash frozen in an isopentane bath cooled to –80C using dry ice. The brain tissue was then embedded in Optimal Cutting Temperature (OCT) compound (Tissue-Tek 62550-12). A cryostat was used to obtain a 10 μm thin section from the hippocampus that was then mounted onto a 10X spatial gene expression slide. Sections were stained with Gfap (Millipore MAB3402), NeuN (Abcam ab177487), and Hoechst 33342 (Invitrogen H1399) before imaging on a Leica Aperio Versa slide scanner. The cDNA libraries were generated at the Gladstone Genomics Core. Libraries were sequenced at the UCSF Center for Advanced Technology on an Illumina NovaSeq 600 on an SP flow cell. Alignment of the sequencing data with spatial data from the Visium slide was completed with the 10X Space Ranger software. The 10X Loupe Browser software was then used to identify four anatomical regions of interest: (1) CA1, (2) CA3, (3) dentate gyrus, and (4) thalamus. RNA capture areas corresponding to each anatomical region were selected for analysis based on their spatial proximity to the anatomical regions and on the expression

of known genetic markers. CA1 genetic markers included *Fibcd1* and *Wfs1*; CA3 markers included *Nrip3* and *Bok*; dentate gyrus markers included *Calb1* and *Prox1*; and the thalamus marker was *Synpo2*. Roughly 15–20 neuronal cell bodies fit into a single capture area for CA1 with roughly 28 capture areas for CA1. Gene expression levels were exported to GraphPad Prism and Microsoft Excel. Gene rankings for hit analysis were established using the fold change score (FCS) and signal-to-noise score (SNS). Equations for these scores are given as:

$$SNS = \frac{\mu_{KO} - \mu_{WT}}{\sigma_{KO} + \sigma_{WT}} \times (-\text{Log}_{10} P)$$

$$FCS = \text{Log}_2 \left(\frac{\mu_{KO}}{\mu_{WT}} \right) \times (-\text{Log}_{10} P)$$

where μ is the average gene expression, σ is the standard deviation, and P is the p value derived from an t test. Genes with a p value <0.05 that also appeared in the top 20% for both scoring metrics were highlighted as differentially expressed genes of interest. The expression levels for these genes of interest were probed in both CA1 and the thalamus to identify a subset of genes with CA1-specific changes. Pathway analysis was done on the differentially expressed genes using the Enrichr webtool^{89–91}

QUANTIFICATION AND STATISTICAL ANALYSIS

All statistical analyses including the n, what n represents, description of error bars, statistical tests used and level of significance are found in the figure legends. Linear mixed modeling and either Welch's ANOVA with Dunnett's T3 multiple comparison test or Welch's t-tests were used to compare WT versus KO animals. Unpaired t-tests were used to compare metabolomics data within a single time point, whereas two-way ANOVA with Sidak's multiple comparison was used for comparisons across timepoints. Mann-Whitney U test was used for electrophysiology data. Visium spatial transcriptomics data of targeted gene sets was analyzed by one or two-way ANOVA with either Sidak's or Tukey's multiple comparisons test. Determination of top differentially expressed genes used a threshold of the top 20% of differentially expressed genes that were ranked according to the defined SNS and FCS metrics. All analyses were performed using GraphPad Prism version 8.4.3 and Microsoft Excel.